

Virbac**Equimax™**

FOR ANIMAL USE ONLY

Equimax™ NF Oral Paste

Reg. No. G3546 (Act 36/1947)

CAUTION**STORAGE INSTRUCTIONS**

Store in a cool dry place.

COMPOSITION

Ivermectin 1,87 % m/m
 Praziquantel 14,03 % m/m

INDICATIONS

Broad spectrum roundworm, tapeworm and bot remedy for horses of all ages, including pregnant mares.

WARNINGS

- Do not use in animals intended for human consumption.
- All pregnant mares must be treated with the usual additional care.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

- Store in a safe place.
- Refrain from smoking and eating whilst handling the product.
- Wash hands after use.

DIRECTIONS FOR USE - Use only as directed

Equimax is administered orally by means of a disposable syringe at a dose rate of 200 µg of ivermectin and 1,5 mg of praziquantel per kg of body mass (1,07 g of oral paste per 100 kg body weight). Each mass marking on the syringe plunger delivers enough paste to treat 100 kg body mass.

Weight	Dosage	Weight	Dosage
Up to 100 kg	1,070 g	401 - 450 kg	4,815 g
101 - 150 kg	1,605 g	451 - 500 kg	5,350 g
151 - 200 kg	2,140 g	501 - 550 kg	5,885 g
201 - 250 kg	2,675 g	551 - 600 kg	6,420 g
251 - 300 kg	3,210 g	601 - 650 kg	6,955 g
301 - 350 kg	3,745 g	651 - 700 kg	7,490 g
351 - 400 kg	4,280 g		

EFFICACY

EQUIMAX broad spectrum parasiticide for horses provides effective control for all major worms of horses including:

Large strongyles	<i>Strongylus vulgaris</i> (adult and arterial larvae) - bloodworm
	<i>Strongylus edentatus</i> (adult and encisted larvae / tissue stage)
	<i>Strongylus equinus</i> (adult)
	<i>Triodontophorus</i> (adult)
Small Strongyles:	<i>Cyathostomum</i> spp. (adults and 4 th stage larvae)
	<i>Cylicocyclus</i> spp. (adults and 4 th stage larvae)
	<i>Cylicostephanus</i> spp. (adults and 4 th stage larvae)

	<i>Cylicodontophorus</i> spp. (adults and 4 th stage larvae)
	<i>Gyalocephalus</i> spp. (adults and 4 th stage larvae)
Ascarids:	
	<i>Parascaris equorum</i> (adults and 3 rd and 4 th stage larvae)
Pinworm:	
	<i>Oxyuris equi</i> (adults and 3 rd and 4 th stage larvae)
Hairworm:	
	<i>Trichostrongylus axei</i> (adults)
Intestinal threadworms	
	<i>Strongyloides westeri</i> (adults)
Large-mouthed stomach worms	
	<i>Habronema muscae</i> (adults)
	<i>Draschia</i> spp. (3 rd stage larvae)
Lungworms	
	<i>Dictyocaulus armfieldi</i> (adults and 4 th stage larvae)
Cestodes	
	<i>Anoplocephala</i> spp.
Bots	
	<i>Gasterophilus</i> spp. (oral and gastric stages)

Equimax is also effective for the control of skin lesions caused by *Habronema* spp. and *Draschia* spp. cutaneous larvae (summer sores) and *Onchocerca* spp. microfilariae (cutaneous onchocerciasis).

ADMINISTRATION

Insert the syringe in the gap between the front and back teeth and deliver paste onto the back of the tongue. Immediately raise the horse's head for a few seconds after dosing to ensure swallowing of the paste. Before administration, ensure that the horse does not have food in its mouth. The body mass of the horse should be determined accurately to maximise the use of the paste.

Suggested Worm Control Programme

All horses should be included in a regular worm control programme.

Mares: Regular dosing at 6 - 8 week intervals on pastures during summer and autumn is recommended.

Stallions: Treat at least twice a year.

Foals: Regular strategic treatment at 6 - 8 week intervals, commencing at approximately two months of age, is recommended.

Bots: Bot fly activity increases in the late spring and usually peaks in the autumn with a consequent accumulation of stomach bots carried in the winter. Treatment against bots attached to the stomach wall is essential to remove established burdens in the autumn and winter.

Consult your veterinarian for a worm control programme.

Horses of all ages, including pregnant mares, breeding stallions and foals can be treated with no adverse effects.

PRESENTATION

Disposable syringe containing 7,49 g paste.

REGISTRATION HOLDER

VIRBAC RSA (Pty) Ltd
 (Reg. No. 1990/003743/07)
 Private Bag X115, Halfway House 1685,
 South Africa
 Tel.: +27 12 657-6000

Zambia Reg. No: 359/742 V
 03/21 C13069-30032004

Virbac**Equimax™**

SLEGS VIR DIEREGEBRUIK

Equimax™ NF Orale Pasta

Reg. No. G3546 (Wet 36/1947)

VERSIGTIG**BERGINGSAAHWYSINGS**

Berg in 'n koel droë plek.

SAMESTELLINGIvermektien 1,87 mg m/m
Prasikwantel 14,03 mg m/m**INDIKASIES**

Breëspektrum rondewurm-, lintwurm- en papiemiddel vir perde van alle ouderdomme, insluitende dragtige merries.

WAARSKUWINGS

- Moenie in diere gebruik wat vir menslike verbruik bestem is nie.
- Alle dragtige merries moet met die gewone bykomende sorg hanteer word.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel obreedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

- Berg in 'n veilige plek.
- Moenie gedurende hantering rook of eet nie.
- Was hande deeglik na gebruik.

GEBRUIKSAANWYSINGS - Gebruik slegs soos aangedui

Equimax word mondelings met behulp van 'n wegdoenbare spuit teen 'n dosis van 200 µg ivermektien en 1,5 mg of prasikwantel per kg of liggaamsmassa (1,07 g orale pasta per 100 kg liggaamsmassa) toegedien. Elke verstelling lewer genoegsame pasta om 100 kg liggaamsmassa te behandel.

Massa	Dosis	Massa	Dosis
Tot 100 kg	1,070 g	401 - 450 kg	4,815 g
101 - 150 kg	1,605 g	451 - 500 kg	5,350 g
151 - 200 kg	2,140 g	501 - 550 kg	5,885 g
201 - 250 kg	2,675 g	551 - 600 kg	6,420 g
251 - 300 kg	3,210 g	601 - 650 kg	6,955 g
301 - 350 kg	3,745 g	651 - 700 kg	7,490 g
351 - 400 kg	4,280 g		

DOELTREFFENDHEID

Equimax is 'n breëspektrum wurmmiddel vir perde wat al die belangrike perdwurms hoogsdoeltreffend beheer:

Groot strongilide	
	<i>Strongylus vulgaris</i> (volwassenes en arteriele larwes) - bloedwurm
	<i>Strongylus edentatus</i> (volwassenes en geïnsisteerde larwes / weefselstadium)
	<i>Strongylus equinus</i> (volwassenes)
	<i>Triodontophorus</i> (volwassenes)
Klien strongilide	
	<i>Cyathostomum</i> spp. (volwassenes en 4 ^{de} stadium larwes)
	<i>Cylicocyclus</i> spp. (volwassenes en 4 ^{de} stadium larwes)

	<i>Cylicostephanus</i> spp. (volwassenes en 4 ^{de} stadium larwes)
	<i>Cylicodontophorus</i> spp. (volwassenes en 4 ^{de} stadium larwes)
	<i>Gyaloccephalus</i> spp. (volwassenes en 4 ^{de} stadium larwes)
Spelwurm	
	<i>Parascaris equorum</i> (volwassenes en 3 ^{de} en 4 ^{de} stadium larwes)
Sambokwurm	
	<i>Oxyuris equi</i> (volwassenes en 3 ^{de} en 4 ^{de} stadium larwes)
Haarwurm	
	<i>Trichostrongylus axei</i> (volwassenes)
Draadwurms	
	<i>Strongyloides westeri</i> (volwassenes)
Grootbekmaagwurms	
	<i>Habronema muscae</i> (volwassenes)
	<i>Draschia</i> spp. (3 ^{de} stadium larwes)
Longwurms	
	<i>Dictyocaulus amfeldi</i> (volwassenes en 4 ^{de} stadium larwes)
Lintwurms	
	<i>Anoplocephala</i> spp.
Papies	
	<i>Gasterophilus</i> spp. (orale en maagstadiums)

Equimax is ook effektief vir die beheer van velletsels wat deur *Habronema* spp. en *Draschia* spp. onderhuidse larwes (somersere) en *Onchocerca* spp. mikrofilariëe (kutane onchoserkiase) veroorsaak word.

TOEDIENING

Plaas die spuit in die gaping tussen voor- en agtertand en plaas die pasta agter op die tong. Lig die perd se kop onmiddellik op na dosering vir 'n paar sekondes om seker te maak dat die middel gesluk word. Voor behandeling, maak seker dat die perd nie voer in die bek het nie. Die liggaamsmassa van die perd moet akkuraat bepaal word om die maksimum werking van die pasta te verseker.

Voorgestelde Program vir Wurmbeheer

Alle perde behoort volgens 'n gereelde wurmbeheer-program gedoseer te word.

Merries: Doseer gereeld met 6 - 8 week tussenposes op weidings gedurende die somer en herfs.

Hingste: Doseer te minstens 2 keer jaarlik.

Vullens: Gereelde strategiese doserings met 6 - 8 week tussenposes word aanbeveel. Begin wanneer die vul omtrent twee maande oud is.

Papies: In die laat lente is die papie-vlieg op sy bedrywigste en bereik sy piek in die herfs, sodoende word papies die winter ingedra. Behandeling teen die papie wat aan die maagwand geheg is, is noodsaaklik om gevestigde besmettings in die herfs en winter te verwyder.

Raadpleeg u veearts om 'n wurmbeheerprogram te bekom.

Perde van alle ouderdomsgroepe, insluitende dragtige merries, teelhingste, en vullens kan sonder enige nadelige gevolge behandel word.

AANBIEDING

Wegdoenbare spuite met 7,49 g pasta.

REGISTRASIEHOUER

VIRBAC RSA (Pty) Ltd, (Reg. No. 1990/003743/07)
Private Bag X115, Halfway House 1685, Suid Afrika
Tel.: +27 12 657-6000

Zambia Reg. No: 359/742 V
03/21 CI3069-30032004

EQUIMAX NF ORAL PASTE 03/21 CI3069-30032004

Approved text (Act 36/1974) or signed off internally (Act 101/1965) and Registration certificate to accompany all checklist

Regulatory Pharmacist / Officer:

- Proof reading and mark-up of Editor's Copy (revision of / change to approved artwork) utilising approved text as reference.
- Proof reading (new artwork) utilising approved text as reference.
- Perform comprehensive check according to requirements of Act 101 / Act 36 / other regulatory bodies, in conjunction with the artwork checklist.
- Reviews Artwork Check List for correctness in line with current legislation and make changes accordingly.
- Read complete Label/Carton/Package Insert for correctness including language, spelling, text, graphics and legals and the Artwork Checklist.

APPROVED

DATE

Product Managers / Specialist/ Business Unit Managers:

- Check artwork layout and marketing images, colours, spelling errors and Product information.
- Check mock-ups/dummy packs
- Provide information to complete Basic Data Sheet for new products
- Proof reading within 3 business days
- Read complete Label/Carton/Package Insert for correctness including language, spelling, text, graphics and legals and the Artwork Checklist.

APPROVED

DATE

Operations Manager/Production Manager:

- Fitness for use
- Compatibility with equipment
- Variable print information: Lot/Expiry/Date of Manufacture
- Presence of pack size
- Presence of barcodes
- Read complete Label/Carton/Package Insert for correctness including language, spelling, text, graphics and legals and the Artwork Checklist.

APPROVED

DATE

Responsible Pharmacist / Regulatory Affairs Manager:

- Approves Master Specification Sheet
- Read complete Label/Carton/Package Insert for correctness including language, spelling, text, graphics and legals and the Artwork Checklist.

APPROVED

DATE
