



Prazifen® Tablets

For Animal Use Only

Prazifen® Tablets

Reg. No. G2602 (Act 36/1947)

②+⑦†

Treatment for roundworm, hookworm, whip-worm and tapeworm infestation in dogs and cats

STORAGE INSTRUCTIONS

Store at room temperature below 25 °C

COMPOSITION

Each tablet contains:

Praziquantel 50 mg

Fenbendazole 500 mg

WARNINGS

- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

PRECAUTIONS

In cases of overdosage, reactions of discomfort, vomiting and diarrhoea may occur. If these symptoms are observed, consult your veterinarian. Treatment is symptomatic and supportive.

EFFICACY

For the treatment of mixed infections with roundworm, hookworm and tapeworm caused by:

Roundworms - *Toxocara canis*
Toxascaris leonina
Hookworms - *Ancylostoma caninum*
Whipworm - *Trichuris vulpis*
Tapeworms - *Echinococcus* spp.
Dipylidium caninum
Taenia spp.

DIRECTIONS FOR USE

Use only as directed

Tablets to be administered orally. The recommended dosage is 5 mg praziquantel and 50 mg fenbendazole per kg body mass, i.e. one tablet per 10 kg body mass.

To ease administration, tablets can be mixed with a small portion of food. Dietary measures or fasting are not necessary. To ease dispensing to clients, the tablets are scored; give a quarter of tablet per 2,5 kg body mass.

PRESENTATION

Blister pack of 12 tablets.

Cartons containing 96 tablets.

*Benzimidazoles + isoquinolines

Registration holder:

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)

Private Bag X115

Halfway House

1685

South Africa

Tel: +27 12 657 000

Namibia Reg. No. V02/18.1.8/20 NSO

39311250 08/19 C12222-20060516



Prazifen® Tablets

Slegs Vir Dieregebruik
Prazifen® Tablets
Reg. No. G2602 (Wet 36/1947)

**Behandeling van rondewurm-,
haakwurm-, sambokwurm- en
lintwurmbesmettings by honde en
katte**

BERGINGSAAWYSINGS

Berg teen kamertemperatuur benede
25 °C

SAMESTELLING

Elke tablet bevat:

Prasikwantel 50 mg
Fenbendasool 500 mg

WAARSKUWINGS

- Hou buite bereik van kinders, oningeligte persone en diere.
- Alhoewel hierdie middel breedvoerig onder 'n wye verskeidenheid van toestande getoets is, mag dit faal as gevolg van verskeie redes. Indien dit vermoed word, raadpleeg 'n veearts en verwittig die registrasiehouer.

VOORSORGMATREËLS

In gevalle van oordosering is reaksies soos ongemaklikheid, braking en diarree waargeneem. As die simptome waargeneem word, raadpleeg u veearts. Behandeling is simptomaties en ondersteunend.

DOELTREFFENDHEID

Vir die behandeling van gemengde besmettings met rondewurms, haakwurms en lintwurms veroorsaak deur:

②+⑦†

Rondewurms - *Toxocara canis*
 - *Toxascaris leonina*
Haakwurms - *Ancylostoma caninum*
Sambokwurm - *Trichuris vulpis*
Lintwurms - *Echinococcus* spp.
 - *Dipylidium caninum*
 - *Taenia* spp.

GEBRUIKSAANWYSINGS

Gebruik slegs soos aangedui

Tablette moet mondelings toegedien word. Die aanbevole dosis is 5 mg prasikwantel en 50 mg fenbendasool per kg liggaamsmassa, d.i. een tablet per 10 kg liggaamsmassa.

Om toediening te vergemaklik, kan tablette met 'n klein porsie kos gemeng word.

Geen dieetaanpassing of uithongering is nodig nie. Om reseptering aan kliënte te vergemaklik, is die tablette met 'n kruis gemerk; gee 'n kwart tablet per 2,5 kg liggaamsmassa.

AANBIEDING

Stulpverpakking van 12 tablette.
Kartonhouers met 96 tablette.

†Bensimidazole + isokolinole

Registrasiehouer:

VIRBAC RSA (Pty) Ltd
(Reg. No. 1990/003743/07)
Privaatsak X115, Halfway House 1685,
Suid-Afrika
Tel: +27 12 657 6000

Namibia Reg. No. V02/18.1.8/20 **NSO**

39311250 08/19 CI2222-20060516