



# Help them bounce back.

A palatable solution to speed up recovery and recuperation in dogs and cats.



NUTRIBOUND is indicated for Peri-surgery conditions, lack of appetite as well as fatigue and low vitality in dogs and cats.







Shaping the future of animal health




# Targeted, balanced and fortified nutrition for recuperation support backed by science



## Liquid Form

-  Poured on food
-  With water
-  On its own
-  Tube or syringe feeding

## Optimal Formula

-  Rich in essential nutrients, Omega-3 and Omega-6, prebiotics and vitamins

## Highly Palatable

-  Good patient compliance
-  Supports healthy eating and drinking

## Key Nutrients Designed to support recuperation

**Zinc, Arginine, Prebiotics**  
Supports immune function <sup>(2-6)</sup>

**Glutamine**  
Supports Gastrointestinal Tract integrity <sup>(7)</sup>

**Zinc, Arginine, Omega 3 and Omega 6**  
Supports skin health <sup>(8-10)</sup>

**Taurine and E Vitamins<sup>6</sup>**  
Antioxidants <sup>(11-12)</sup>

**Inulin and Fructooligosaccharides**  
Supports Intestinal Microbial Balance <sup>(13-15)</sup>



# Highly palatable recuperation and recovery supplement



**95% prehension rate in cats**

**92%**<sup>(1)</sup> of the total consumption\*



**100% prehension rate in dogs**

**86%**<sup>(1)</sup> of the total consumption\*

\* Total consumption being deemed as a consumption of more than 95% of the product. % among animals which already prehended the product.


## Published Data



**STIMULATION OF WATER INTAKE**

With NUTRIBOUND, cats drink

**46%**<sup>(1)</sup> more




**WATER INTAKE**



**STIMULATION OF FOOD INTAKE**

With NUTRIBOUND, cats eat

**63%**<sup>(1)</sup> more



**FOOD CONSUMPTION**



# When should NUTRIBOUND formula be used?



## Peri-surgery conditions

Liquid feed can be given directly after the surgery in order to help maintain the integrity of the digestive tracts.

Optimal functioning of the immune system is crucial after surgery for faster and more effective recuperation.

- 1 Promoting normal intestinal absorption
- 2 Delivering fluids
- 3 Providing essential nutrients for recuperation
- 4 Providing prebiotic fibers to stimulate intestinal function



## Lack of appetite

High palatability and liquid formulation will help improve appetite.

Prolonged anorexia will lead to an atrophy of the intestinal villi, which can severely impair the absorption of nutrients into the blood.



## Fatigue & low vitality

Because NUTRIBOUND formula is highly palatable, it stimulates the animal to drink and eat whilst providing the essential nutrients for recuperation.



# Early and progressive re-feeding

## 1 START-UP NUTRIBOUND



**For optimal recuperation, stimulating drinking and eating within the first 48 hours is key to a pet's recovery.**

**From the first hours, pets can be given NUTRIBOUND in order to promote hydration and provide essential nutrients.**

## 2 FOLLOW-UP NUTRIBOUND + High energy feed



**Kcal  
++**

**After 48 hours, introduction of a complete diet is recommended.**

To increase tolerance, small and repeated intakes of high energy and easily digestible nutrients are recommended in addition to NUTRIBOUND.

## 3 TRANSITION NUTRIBOUND + Usual diet



NUTRIBOUND can be given with the usual diet to **help pets to return to their normal alimentary habits.**



Helps stimulate eating and drinking.





## NUTRIBOUND®

For animal use only.

For full prescribing details refer to package insert.

### NUTRIBOUND for Cats.

Composition: Meat and animal derivatives (4,07 %), derivatives of vegetable origin (6,70%), oils and fats (1,50 %), minerals (0,62 %), chicory inulin (0,3 %), fructo-oligosaccharides (0,11%).

Reg. No: V29044 (Act 36/1947), Namibia Reg. No: N-FF 4237

### NUTRIBOUND for Dogs.

Composition: Meat and animal derivatives (4,74 %), derivatives of vegetable origin (6,14 %), oils and fats (1,29 %), chicory inulin (0,3 %), fructo-oligosaccharides (0,11 %), minerals (0,69 %).

Reg. No: V29045 (Act 36/1947), Namibia Reg. No: N-FF 4238



Virbac RSA (Pty) Ltd (Reg. No. 1990/003743/07)  
Private Bag X115, Halfway House 1685  
Tel: (012) 657-6000 • Fax: (012) 657-6067

© 08/2018 Virbac all rights reserved

1. Chaix G. 2015, Palatability and impact on food and water intake of Nutribound, a new liquid nutritional supplement for dogs and cats - Journal of Veterinary Science & Medicine. 2. Barbul A, Sisto DA, Wasserkrug HL, Efron G. Arginine stimulates lymphocyte immune response in healthy human beings. Surgery. 1981; 90(2): 244-51. 3. Dupertuis YM, Raguso CA, Richard C. Basics in clinical nutrition, Nutrients which in uence immunity: clinical and experimental data. Eur e-Journ Clin Nutr Metabol. 2009; 4: e7-e9. 4. Field J, Johnson R, Schley D. Nutrients and their role in host resistance to infection. Journal of Leukocyte Biology 2002; 71 (1):16-32. 5. Mocchegiani E, Muzzioli M, Cipriano C, Giacconi R. Zinc, T-cell pathways, aging: role of metallothioneins. Mech Ageing Dev. 1998; 106 (1-2): 183-204. 6. Meyer D. Prebiotic dietary bras and the immune system. Agro Food industry hi-tech. 2008;19 (3): 11-15. 7. Van der Hulst RR, van Kreel BK, von Meyenfeldt MF, Brummer RJ, Arends JW, Deutz NE, Soeters PB. Glutamine and the preservation of gut integrity. Lancet. 1993; 29 (8857): 1363-1365. 8. Mackay D, Miller A.L. Nutritional Support for Wound Healing. Alternative Medicine Review. 2003; 8 (4): 359-377. 9. Lansdown AB, Mirastschijski U, Stubbs N, Scanlon E, Agren MS. Zinc in wound healing: theoretical, experimental, and clinical aspects. Wound Repair Regen. 2007 ; 15(1): 2-16. 10. Campbell. Clinical Use of Fatty Acid Supplements in Dogs. Veterinary Dermatology. 1993; 4 (4): 167-173. 11. Traber, Maret G; Stevens, Jan F. Free Radical Biology and Medicine - Vitamins C and E: Bene cial effects from a mechanistic perspective. Free Radical Biology and Medicine 2011; 51 (5): 1000-1013. 12. Grimble RF. Basics in clinical nutrition: immunonutrition. Nutrients which in uence immunity: effect and mechanism of action. Eur e-Journ Clin Nutr Metabol. 2009; 4: e10-e13. 13. Flickinger EA, Fahey GC Jr. Pet food and feed applications of inulin, oligofructose and other oligosaccharides. Br J Nutr. 2002; 87 (2): 297-300. 14. Cummings JH, MacFarlane GT. Gastrointestinal effects of prebiotics. Br J Nutr 2002; 87 (2): 145-151. 15. Verdonk JM, Shim SB, van Leeuwen P, Versteegen MW. Application of inulin-type fructans in animal feed and pet food. Br J Nutr. 2005; 93 (1): 125-38